



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

**NATIONAL  
SENIOR CERTIFICATE/  
NASIONALE SENIOR  
SERTIFIKAAT**

**GRADE 12/GRAAD 12**

**MATHEMATICS P1/WISKUNDE VI  
NOVEMBER 2021  
MARKING GUIDELINES/NASIENRIGLYNE**

**MARKS/PUNTE: 150**

**These marking guidelines consist of 16 pages.  
*Hierdie nasienriglyne bestaan uit 16 bladsye.***


**NOTE:**

- If a candidate answers a question TWICE, only mark the FIRST attempt.
- Consistent accuracy applies in all aspects of the marking guidelines.

**LET WEL:**

- Indien 'n kandidaat 'n vraag TWEE keer beantwoord, sien slegs die EERSTE poging na.
- Volgehoue akkuraatheid is op ALLE aspekte van die nasienriglyne van toepassing.

**QUESTION/VRAAG 1**

1.1.1	$x^2 - 2x - 24 = 0$ $(x - 6)(x + 4) = 0$ $x = 6$ or $x = -4$	✓ factors ✓ $x = 6$ ✓ $x = -4$ (3)
1.1.2	$2x^2 - 3x - 3 = 0$ $x = \frac{3 \pm \sqrt{(-3)^2 - 4(2)(-3)}}{2(2)}$ $x = \frac{3 \pm \sqrt{33}}{4}$ $x = 2,19$ or $x = -0,69$	✓ substitution into the correct formula ✓ $x = 2,19$ ✓ $x = -0,69$ (3)
1.1.3	$x^2 + 5x \leq -4$ $x^2 + 5x + 4 \leq 0$ $(x + 4)(x + 1) \leq 0$ Critical values: $x = -4$ or $x = -1$  $-4 \leq x \leq -1$ <b>OR/OF</b> $x \in [-4 ; -1]$	✓ standard form ✓ critical values ✓ ✓ answer (4)
1.1.4	$\sqrt{x + 28} = 2 - x$ $(\sqrt{x + 28})^2 = (2 - x)^2$ $x + 28 = 4 - 4x + x^2$ $x^2 - 5x - 24 = 0$ $(x - 8)(x + 3) = 0$ $x \neq 8$ or $x = -3$	✓ squaring both sides ✓ standard form ✓ factors ✓ answers with selection (4)

<p>1.2</p>	$2y = 3 + x$ $x = 2y - 3 \quad \dots (1)$ $2xy + 7 = x^2 + 4y^2 \quad \dots (2)$ $2y(2y - 3) + 7 = (2y - 3)^2 + 4y^2$ $4y^2 - 6y + 7 = 4y^2 - 12y + 9 + 4y^2$ $4y^2 - 6y + 2 = 0$ $2y^2 - 3y + 1 = 0$ $(2y - 1)(y - 1) = 0$ $y = \frac{1}{2} \text{ or } y = 1$ $x = -2 \text{ or } x = -1$ <p><b>OR/OF</b></p> $2y = 3 + x$ $y = \frac{3}{2} + \frac{x}{2} \quad \dots(1)$ $2xy + 7 = x^2 + 4y^2 \quad \dots (2)$ $2x\left(\frac{3}{2} + \frac{x}{2}\right) + 7 = x^2 + 4\left(\frac{3}{2} + \frac{x}{2}\right)^2$ $3x + x^2 + 7 = x^2 + 9 + 6x + x^2$ $x^2 + 3x + 2 = 0$ $(x + 2)(x + 1) = 0$ $x = -2 \text{ or } x = -1$ $y = \frac{1}{2} \text{ or } y = 1$	<p>✓ equation 1</p> <p>✓ substitution ✓ simplification</p> <p>✓ standard form</p> <p>✓ y – values ✓ x – values</p> <p>(6)</p> <p><b>OR/OF</b></p> <p>✓ equation 1</p> <p>✓ substitution</p> <p>✓ simplification ✓ standard form</p> <p>✓ x – values ✓ y – values</p> <p>(6)</p>
<p>1.3</p>	$\frac{n}{m} = \frac{p}{n}$ $n^2 = mp$ $\Delta = b^2 - 4ac$ $\Delta = n^2 - 4mp, \text{ but } n^2 = mp$ <p><b>OR/OF</b></p> $\Delta = n^2 - 4n^2 \quad \Delta = mp - 4mp$ $\Delta = -3n^2 \quad \Delta = -3mp$ $n^2 > 0 \quad mp > 0$ $\therefore -3n^2 < 0 \quad \therefore -3mp < 0$ <p><math>\therefore \Delta &lt; 0 \Rightarrow x</math> is a non-real number</p>	<p>✓ <math>\frac{n}{m} = \frac{p}{n}</math> ✓ <math>n^2 = mp</math></p> <p>✓ <math>\Delta = -3n^2</math> or <math>-3mp</math></p> <p>✓ <math>\Delta &lt; 0</math></p> <p>(4)</p>
<p><b>[24]</b></p>		

## QUESTION/VRAAG 2

2.1	$\frac{90}{x} = \frac{81}{90}$ $81x = 8100$ $x = 100$ <p><b>OR/OF</b></p> $x = 90 \times \frac{10}{9}$ $x = 100$	$\checkmark \frac{90}{x} = \frac{81}{90}$ $\checkmark \text{ answer} \quad (2)$ <p><b>OR/OF</b></p> $\checkmark \frac{10}{9}$ $\checkmark \text{ answer} \quad (2)$
2.2	$S_n = \frac{a(1-r^n)}{1-r}$ $S_n = \frac{100(1-(0,9)^n)}{1-0,9}$ $S_n = \frac{100(1-(0,9)^n)}{0,1}$ $\therefore S_n = 1\,000(1-(0,9)^n)$	$\checkmark r = 0,9$ $\checkmark \text{ substitution into correct formula} \quad (2)$
2.3	$S_\infty = \frac{a}{1-r}$ $S_\infty = \frac{100}{1-\frac{9}{10}}$ $S_\infty = 1000$ <p><b>OR/OF</b></p> $S_\infty = \lim_{n \rightarrow \infty} [1\,000(1-(0,9)^n)]$ $S_\infty = 1000$	$\checkmark \text{ substitution}$ $\checkmark \text{ answer} \quad (2)$ <p><b>OR/OF</b></p> $\checkmark S_\infty = \lim_{n \rightarrow \infty} [1\,000(1-(0,9)^n)]$ $\checkmark \text{ answer} \quad (2)$
<b>[6]</b>		


**QUESTION 3**

3.1	-82	✓ answer (1)
3.2	<p>-145 ; -122 ; -101 ; ...</p> $\begin{array}{ccc} & \swarrow & \searrow \\ & 23 & 21 \\ & \swarrow & \searrow \\ & -2 & -2 \end{array}$ <p> <math>2a = -2 \quad \therefore a = -1</math>  <math>3a + b = 23 \quad \therefore 3(-1) + b = 23 \quad \therefore b = 26</math>  <math>a + b + c = -145 \quad \therefore -1 + 26 + c = -145 \quad \therefore c = -170</math>  <math>\therefore T_n = -n^2 + 26n - 170</math> </p> <p><b>OR/OF</b></p> <p> <math>2a = -2 \quad \therefore a = -1</math>  <math>c = -145 + (-2) - 23 = -170</math>  <math>\therefore T_n = -n^2 + bn - 170</math>  <math>-145 = -1 + b - 170</math>  <math>b = 26</math>  <math>\therefore T_n = -n^2 + 26n - 170</math> </p>	<p> <math>\checkmark 2a = -2</math>  <math>\checkmark 3(-1) + b = 23</math>  <math>\checkmark -1 + 26 + c = -145</math> </p> <p>(3)</p> <p><b>OR/OF</b></p> <p> <math>\checkmark 2a = -2</math>  <math>\checkmark c = -145 + (-2) - 23</math> </p> <p> <math>\checkmark -145 = -1 + b - 170</math> </p> <p>(3)</p>
3.3	<p> <math>T_n = bn + c</math>    or    <math>T_n = a + (n - 1)d</math>  <math>T_n = -2n + 25</math>    = <math>23 + (n - 1)(-2)</math>  <math>-2n + 25 = -121</math>    = <math>25 - 2n</math>  <math>-2n = -146</math>  <math>n = 73</math>                      Between <math>T_{73}</math> and <math>T_{74}</math> </p> <p><b>OR/OF</b></p> <p> <math>T_{n+1} - T_n = -(n + 1)^2 + 26(n + 1) - 170 - (-n^2 + 26n - 170)</math>  <math>-121 = -2n + 25</math>  <math>n = 73</math>                      Between <math>T_{73}</math> and <math>T_{74}</math> </p>	<p> <math>\checkmark T_n = -2n + 25</math>  <math>\checkmark T_n = -121</math>  <math>\checkmark n = 73</math>  <math>\checkmark</math> answer                 </p> <p>(4)</p> <p><b>OR/OF</b></p> <p> <math>\checkmark T_n = -2n + 25</math>  <math>\checkmark T_n = -121</math>  <math>\checkmark n = 73</math>  <math>\checkmark</math> answer                 </p> <p>(4)</p>
3.4	<p> <math>n = \frac{-b}{2a} = \frac{-26}{2(1)} = 13</math>  <math>T_{13} = -1</math>  <math>\therefore</math> add 2                 </p> <p><b>OR/OF</b></p> <p> <math>T'_n = -2n + 26 = 0</math>  <math>n = 13</math>  <math>T_{13} = -(13)^2 + 26(13) - 170 = -1</math>  <math>\therefore</math> add 2                 </p>	<p> <math>\checkmark 13</math>  <math>\checkmark T_{13} = -1</math>  <math>\checkmark</math> add 2                 </p> <p>(3)</p> <p><b>OR/OF</b></p> <p> <math>\checkmark 13</math>  <math>\checkmark T_{13} = -1</math>  <math>\checkmark</math> add 2                 </p> <p>(3)</p>

**QUESTION/VRAAG 4**

4.1	$a = 5$ and/en $d = 2$ $T_{51} = 5 + (51 - 1)(2)$ $= 105$	✓ $a$ and $d$ ✓ substitution into correct formula ✓ answer (3)
4.2	$S_n = \frac{n}{2}[2a + (n - 1)d]$ $S_{51} = \frac{51}{2}[2(5) + (51 - 1)2]$ or/of $S_{51} = \frac{51}{2}[5 + 105]$ $= 2\ 805$ $= 2\ 805$	✓ substitution into correct formula ✓ answer (2)
4.3	$\sum_{n=1}^{5\ 000} (2n + 3) = 5 + 7 + 9 + \dots + 10\ 003$	✓ expansion (1)
4.4	$T_1 = -3$ $T_{4\ 999} = -2(4\ 999) - 1 = -9\ 999$  $\therefore \sum_{n=1}^{5\ 000} (2n + 3) + \sum_{n=1}^{4\ 999} (-2n - 1)$ $= (5 + 7 + 9 + \dots + 9\ 999 + 10\ 001 + 10\ 003) +$ $(-3 - 5 - 7 - 9 - \dots - 9\ 999)$ $= 10\ 001 + 10\ 003 - 3$ $= 20\ 001$  <b>OR/OF</b>  $S_{4\ 999} = \frac{4\ 999}{2}[2(-3) + (4\ 999 - 1)(-2)] = -24\ 999\ 999$  $S_{5\ 000} = \frac{5\ 000}{2}((2)(5) + (5\ 000 - 1)(2)) = 25\ 020\ 000$  $\sum_{n=1}^{5\ 000} (2n + 3) + \sum_{n=1}^{4\ 999} (-2n - 1) = 25\ 020\ 000 - 24\ 999\ 999$ $= 20\ 001$	✓ $T_1 = -3$ ✓ $T_{4\ 999} = -9\ 999$  ✓ both expansions  ✓ answer (A) (4)  <b>OR/OF</b>  ✓ $T_1 = -3$ ✓ $S_{4\ 999} = -24\ 999\ 999$  ✓ $S_{5\ 000} = 25\ 020\ 000$  ✓ answer (A) (4)
		<b>[10]</b>

**QUESTION/VRAAG 5**

5.1	$x = 3$ $y = 2$	✓ $x = 3$ ✓ $y = 2$ (2)
5.2	$x \in R, x \neq 3$  <b>OR/OF</b>  $x \in (-\infty ; 3) \cup (3 ; \infty)$  <b>OR/OF</b>  $x < 3$ or $x > 3$	✓ answer (1)  <b>OR/OF</b>  ✓ answer (1)  <b>OR/OF</b>  ✓ answer (1)
5.3	$0 = \frac{-1}{x-3} + 2$ $-2x + 6 = -1$ $x = \frac{7}{2}$ x-int: $\left(\frac{7}{2} ; 0\right)$	✓ $y = 0$  ✓ answer (2)
5.4	y-int: $\left(0 ; \frac{7}{3}\right)$	✓ $x = 0$ ✓ $\frac{7}{3}$ (2)
5.5		✓ asymptotes ✓ intercepts with the axes ✓ shape (3)
		<b>[10]</b>

**QUESTION/VRAAG 6**

6.1	$f(x) = \log_4 x$ $2 = \log_4 k$ $4^2 = k$ $\therefore k = 16$	✓ substitution of $(k ; 2)$  ✓ answer  (2)
6.2	$-1 = \log_4 x \quad \therefore x = \frac{1}{4}$  $\frac{1}{4} \leq x \leq 16$ or/of $x \in \left[ \frac{1}{4} ; 16 \right]$	✓ $x = \frac{1}{4}$  ✓ answer  (2)
6.3	$f(x) = \log_4 x$ $y = \log_4 x$ $x = \log_4 y$ $y = 4^x$	✓ swopping $x$ and $y$  ✓ answer  (2)
6.4	$x < 0$  <b>OR/OF</b>  $x \in (-\infty ; 0)$	✓✓ answer  <b>OR/OF</b>  ✓✓ answer  (2)  (2)
		<b>[8]</b>



**QUESTION 7**

7.1	$B(-4 ; 0)$ $D(6 ; 0)$	$\checkmark B(-4 ; 0)$ $\checkmark D(6 ; 0)$ (2)
7.2	$f(x) = x^2 - 2x - 24$ $x_{tp} = \frac{-b}{2a}$ <b>OR/OF</b> $2x - 2 = 0$ <b>OR/OF</b> $x = \frac{-4+6}{2}$ $x = \frac{-(-2)}{2(1)}$ $\therefore x_{tp} = 1$ $y_{tp} = f(1)$ $= 1^2 - 2(1) - 24$ $= -25$ $C(1 ; -25)$	$\checkmark x_{tp} = 1$  $\checkmark y_{tp} = -25$ (2)
7.3	$y \geq -25$  <b>OR/OF</b> $y \in [-25 ; \infty)$	$\checkmark$ answer (1)  <b>OR/OF</b> $\checkmark$ answer (1)
7.4.1	$m_{AE} = \tan 14,04^\circ = 0,25 = \frac{1}{4}$	$\checkmark$ answer (1)
7.4.2	$m_{\text{tang}} = -4$ $f'(x) = 2x - 2$  $2x - 2 = -4$ $x_T = -1$ $y_T = -21$	$\checkmark m_{\text{tang}} = -4$ $\checkmark f'(x) = 2x - 2$  $\checkmark$ equating $\checkmark x_T = -1$ $\checkmark y_T = -21$ (5)
7.5	$m_{\text{line}} = \frac{1}{4}$ $y + 9 = \frac{1}{4}(x + 3)$ <b>OR/OF</b> $-9 = \frac{1}{4}(-3) + c$ $y + 9 = \frac{1}{4}x + \frac{3}{4}$ $c = -\frac{33}{4} = -8,25$ $y = \frac{1}{4}x - \frac{33}{4}$ <b>OR/OF</b> $y = 0,25x - 8,25$  $x^2 - 2x - 24 = \frac{1}{4}x - \frac{33}{4}$ $4x^2 - 8x - 96 = x - 33$ $4x^2 - 9x - 63 = 0$ $(4x - 21)(x + 3) = 0$ $\therefore x = \frac{21}{4} = 5,25$ or $x \neq -3$	$\checkmark m_{\text{line}} = \frac{1}{4}$  $\checkmark$ substitution $m$ and $K(-3 ; -9)$  $\checkmark y = \frac{1}{4}x - \frac{33}{4}$  $\checkmark$ equating $\checkmark$ standard form  $\checkmark$ answer with selection (6)
		<b>[17]</b>

**QUESTION/VRAAG 8**

<p>8.1</p>	$A = P(1 - i)^n$ $A = 980\,000(1 - 0,092)^7$ $A = R498\,685,82$	<p>✓ correct formula                  ✓ substitution                  ✓ answer (A) (3)</p>
<p>8.2</p>	$A = P(1 + i)^n$ $116\,253,50 = 75\,000 \left(1 + \frac{0,068}{4}\right)^{4n}$ $1,550\,046\,667 = (1,017)^{4n}$ $\log(1,550\,046\,667) = 4n \log(1,017)$ $4n = \frac{\log(1,550\,046\,667)}{\log(1,017)} \text{ or } 4n = \log_{1,017}(1,550\,046\,667)$ $4n = 25,99 \dots$ $n = 6,50 \text{ years}$	<p>✓ <math>\frac{0,068}{4}</math>                  ✓ substitution in correct formula                  ✓ correct use of logs                  ✓ answer (4)</p>
<p>8.3.1</p>	$F = \frac{x[(1 + i)^n - 1]}{i}$ $450\,000 = \frac{x \left[ \left(1 + \frac{0,0835}{12}\right)^{60} - 1 \right]}{\frac{0,0835}{12}}$ $x = R6\,068,69$	<p>✓ <math>i = \frac{0,0835}{12}</math>                  ✓ substitution into correct formula                  ✓ answer (3)</p>
<p>8.3.2(a)</p>	$P = \frac{x[1 - (1 + i)^{-n}]}{i}$ $P = \frac{11\,058,85 \left[ 1 - \left(1 + \frac{0,12}{12}\right)^{-4 \times 12} \right]}{\frac{0,12}{12}}$ $P = R419\,948,32$ <p><b>OR/OF</b></p> <p>Balance = A - F</p> $= P(1 + i)^n - \frac{x[(1 + i)^n - 1]}{i}$ $= 1\,050\,000 \left(1 + \frac{0,12}{12}\right)^{12 \times 21} - \frac{11\,058,85 \left[ \left(1 + \frac{0,12}{12}\right)^{12 \times 21} - 1 \right]}{\frac{0,12}{12}}$ $= R12\,887\,702,20 - R12\,467\,749,81$ $= R419\,952,39$	<p>✓ <math>n = 48</math> in P-formula                  ✓ substitution into correct formula                  ✓ answer (A) (3)</p> <p><b>OR/OF</b></p> <p>✓ <math>n = 252</math> in both formulae                  ✓ subst into correct formulae                  ✓ answer (A) (3)</p>

<p>8.3.2(b)</p>	<p>Total paid = <math>11\,058,85 \times 21 \times 12 = 2\,786\,830,20</math>                  Loan Paid = <math>1\,050\,000 - 419\,948,32 = 630\,051,68</math>                  Interest paid = <math>2\,786\,830,20 - 630\,051,68</math>                                    = R2 156 778,52</p> <p><b>OR/OF</b></p> <p>Total paid = <math>11\,058,85 \times 21 \times 12 = 2\,786\,830,20</math>                  Loan Paid = <math>1\,050\,000 - 419\,952,39 = 630\,047,61</math>                  Interest paid = <math>2\,786\,830,20 - 630\,047,61</math>                                    = R2 156 782,59</p> <p><b>OR/OF</b></p> <p>Interest paid                  = <math>11\,058,85 \times 21 \times 12 - (1\,050\,000 - 419\,948,32)</math>                  = <math>2\,786\,830,20 - 630\,051,68</math>                  = R2 156 778,52</p>	<p>✓ <math>11\,058,85 \times 21 \times 12</math>                  ✓ <math>1\,050\,000 - \text{Balance Outstanding}</math>                  ✓ answer (3)</p> <p><b>OR/OF</b></p> <p>✓ <math>11\,058,85 \times 21 \times 12</math>                  ✓ <math>1\,050\,000 - \text{Balance Outstanding}</math>                  ✓ answer (3)</p> <p><b>OR/OF</b></p> <p>✓ <math>11\,058,85 \times 21 \times 12</math>                  ✓ <math>1\,050\,000 - \text{Balance Outstanding}</math>                  ✓ answer (3)</p>
		<p><b>[16]</b></p>

## QUESTION/VRAAG 9

9.1	$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{2(x+h)^2 - 3(x+h) - (2x^2 - 3x)}{h}$ $f'(x) = \lim_{h \rightarrow 0} \frac{2x^2 + 4xh + 2h^2 - 3x - 3h - 2x^2 + 3x}{h}$ $= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - 3h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 3)}{h}$ $= \lim_{h \rightarrow 0} (4x + 2h - 3)$ $\therefore f'(x) = 4x - 3$ <p><b>OR/OF</b></p> $f(x) = 2x^2 - 3x$ $f(x+h) = 2(x+h)^2 - 3(x+h)$ $f(x+h) = 2x^2 + 4xh + 2h^2 - 3x - 3h$ $f(x+h) - f(x) = 4xh + 2h^2 - 3h$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - 3h}{h}$ $= \lim_{h \rightarrow 0} \frac{h(4x + 2h - 3)}{h}$ $= \lim_{h \rightarrow 0} (4x + 2h - 3)$ $\therefore f'(x) = 4x - 3$	<p>✓ substitution</p> <p>✓ <math>2x^2 + 4xh + 2h^2 - 3x - 3h</math></p> <p>✓ <math>4xh + 2h^2 - 3h</math></p> <p>✓ factorisation</p> <p>✓ answer (5)</p> <p><b>OR/OF</b></p> <p>✓ substitution</p> <p>✓ <math>2x^2 + 4xh + 2h^2 - 3x - 3h</math></p> <p>✓ <math>4xh + 2h^2 - 3h</math></p> <p>✓ factorisation</p> <p>✓ answer (5)</p>
9.2.1	$y = 4x^5 - 6x^4 + 3x$ $\frac{dy}{dx} = 20x^4 - 24x^3 + 3$	<p>✓ <math>20x^4</math></p> <p>✓ <math>-24x^3</math></p> <p>✓ 3 (3)</p>

9.2.2	$D_x \left[ \frac{-\sqrt[3]{x}}{2} + \left( \frac{1}{3x} \right)^2 \right]$ $D_x \left[ \frac{-x^{\frac{1}{3}}}{2} + \frac{x^{-2}}{9} \right]$ $D_x \left[ -\frac{1}{2} x^{\frac{1}{3}} + \frac{1}{9} x^{-2} \right]$ $= -\frac{1}{6} x^{-\frac{2}{3}} - \frac{2x^{-3}}{9}$ $= -\frac{1}{6x^{\frac{2}{3}}} - \frac{2}{9x^3}$	$\checkmark \frac{-x^{\frac{1}{3}}}{2} \quad \checkmark \frac{x^{-2}}{9}$  $\checkmark -\frac{1}{6} x^{-\frac{2}{3}} \quad \checkmark -\frac{2x^{-3}}{9}$  (4)
		<b>[12]</b>

**QUESTION/VRAAG 10**

<p>10.1</p>	<p><math>h(x) = ax^3 + bx^2</math>  <math>h'(x) = 3ax^2 + 2bx</math>  <math>h'(4) = 0</math>  <math>48a + 8b = 0</math>  <math>6a + b = 0 \quad \dots(1)</math></p> <p><math>h(4) = 32</math>  <math>64a + 16b = 32</math>  <math>4a + b = 2 \quad \dots(2)</math></p> <p><math>(1) - (2): 6a + b = 0</math>  <math>4a + b = 2</math></p> <p style="margin-left: 40px;"><math>2a = -2</math>  <math>a = -1</math></p> <p style="margin-left: 40px;"><math>4(-1) + b = 2</math>  <math>b = 6</math></p>	<p>✓ <math>h'(x)</math>                  ✓ <math>h'(4) = 0</math>                  ✓ <math>48a + 8b = 0</math> or <math>6a + b = 0</math></p> <p>✓ <math>h(4) = 32</math>                  ✓ <math>64a + 16b = 32</math> or <math>4a + b = 2</math></p> <p style="text-align: right;">(5)</p>
<p>10.2</p>	<p><math>h(x) = -x^3 + 6x^2</math>  <math>-x^3 + 6x^2 = 0</math>  <math>x^2(-x + 6) = 0</math>  <math>x = 0</math> or <math>x = 6</math>  <math>\therefore A(6; 0)</math></p>	<p>✓ <math>-x^3 + 6x^2 = 0</math>                  ✓ factors</p> <p>✓ <math>A(6; 0)</math></p> <p style="text-align: right;">(3)</p>
<p>10.3.1</p>	<p><math>0 &lt; x &lt; 4</math> or <math>0 \leq x \leq 4</math></p> <p><b>OR/OF</b></p> <p><math>x \in (0; 4)</math> or <math>x \in [0; 4]</math></p>	<p>✓ critical values                  ✓ notation (2)</p> <p><b>OR/OF</b></p> <p>✓ critical values                  ✓ notation (2)</p>
<p>10.3.2</p>	<p><math>x &gt; 2</math></p> <p><b>OR/OF</b></p> <p><math>x \in (2; \infty)</math></p>	<p>✓ 2                  ✓ notation (2)</p> <p><b>OR/OF</b></p> <p>✓ 2                  ✓ notation (2)</p>
<p>10.4</p>	<p><math>f(x) = h(x-1) = -(x-1)^3 + 6(x-1)^2</math>  <math>f(0) = 7</math>  <math>7 &lt; k &lt; 32</math> or <math>k \in (7; 32)</math></p>	<p>✓ <math>k &lt; 32</math>                  ✓ new y-intercept = 7                  ✓ <math>7 &lt; k &lt; 32</math></p> <p style="text-align: right;">(3)</p>
		<p>[15]</p>

**QUESTION/VRAAG 11**

11	$\text{Time} = \frac{20}{x}$ $\text{Cost} = (\text{water cost per hour} \times \text{time}) + (\text{kms} \times \text{R/km})$ $C(x) = 1,6 \times \left(\frac{20}{x}\right) + 20 \left(1,2 + \frac{x}{4000}\right)$ $C(x) = \frac{32}{x} + 24 + \frac{x}{200}$ $C'(x) = -\frac{32}{x^2} + \frac{1}{200} = 0$ $x^2 = 6400$ $x = 80 \text{ km/h}$	$\checkmark \frac{20}{x}$ $\checkmark 1,6 \times \left(\frac{20}{x}\right)$ $\checkmark 20 \left(1,2 + \frac{x}{4000}\right)$ $\checkmark C(x) = \frac{32}{x} + 24 + \frac{x}{200}$ $\checkmark C'(x) = -\frac{32}{x^2} + \frac{1}{200}$ $\checkmark C'(x) = 0$ $\checkmark \text{answer (A)}$
		(7)
		[7]

**QUESTION/VRAAG 12**

12.1.1	No, because $P(A \text{ and } B) \neq 0$	$\checkmark \text{answer and reason}$
		(1)
12.1.2(a)	$P(A \text{ and } B) = 0,3 \quad P(\text{only } B) = 0,2$ $P(A \text{ and } B) = P(A) \times P(B)$ $0,3 = P(A) \times 0,5$ $P(A) = 0,6$ $P(\text{only } A) = 0,3$	$\checkmark P(A \text{ and } B) = P(A) \times P(B)$ $\checkmark 0,5$ $\checkmark P(A) = 0,6$ $\checkmark \text{answer}$
		(4)
12.1.2(b)	<div style="border: 1px solid black; padding: 10px; margin-bottom: 10px;"> </div> $P(\text{not } A \text{ or not } B) = 0,2 + 0,2 + 0,3 = 0,7$ <p><b>OR/OF</b></p> $P(\text{not } A \text{ or not } B) = 1 - P(A \text{ and } B) = 1 - 0,3 = 0,7$ <p><b>OR/OF</b></p> $P(A' \text{ or } B') = P(A') + P(B') - P(A' \text{ and } B')$ $= 0,4 + 0,5 - 0,2 = 0,7$	$\checkmark \text{method}$ $\checkmark \text{answer}$
		(2)
		(2)
		(2)

12.2.1	$P(\text{novel}) = \frac{3}{12} = \frac{1}{4}$				✓ answer (1)
12.2.2	$12! = 479\ 001\ 600$				✓✓ answer (2)
12.2.3	5 (Poetry)	3! (Novels all together)	8! (Arrangements of rest of the books including the novels)	4 (Drama)	✓ $5 \times 4$ ✓ $3! = 6$ ✓ $8!$  $P(\text{start with poetry, end with drama AND all novels together})$ $= \frac{5 \times 3! \times 8! \times 4}{12!}$ $= \frac{1}{99}$
					✓ $\frac{5 \times 3! \times 8! \times 4}{12!} = \frac{1}{99}$ (A) (4)
					[14]

**TOTAL/TOTAAL: 150**